

Answer on Question #73562 – Math – Calculus

Question

Evaluate the integral by converting to polar coordinate integrate 0 to $\sqrt{3}$ double integrate y to $\sqrt{4-y^2}$ dx.dy/(4+ x^2+y^2).

Solution

$$\text{Given } \int_0^{\sqrt{3}} \int_y^{\sqrt{4-y^2}} \frac{dx dy}{4+x^2+y^2}$$

By the definition of the polar coordinates $x = r \cos \theta$, $y = r \sin \theta$, $x^2 + y^2 = r^2$, $dx dy = r dr d\theta$

First of all limits of y are given 0 and $\sqrt{3}$ here $y = r \sin \theta$, when $y = 0$ then $\theta = 0$, $y = \sqrt{3}$

$$\text{then } \theta = \frac{\pi}{3}$$

Similarly here

$$x = r \cos \theta, \quad x = \sqrt{4-y^2}, \quad x^2 + y^2 = r^2, \quad r^2 = 4, \quad r = 2$$

For the region of integration, r varies from 0 to 2 and $\theta = 0$ and $\theta = \frac{\pi}{3}$

$$\int_0^{\frac{\pi}{3}} \int_0^2 \frac{r dr d\theta}{4+r^2}$$

$$= \frac{1}{2} \int_0^{\frac{\pi}{3}} \int_0^2 \frac{2r dr d\theta}{4+r^2} = \frac{1}{2} \int_0^{\frac{\pi}{3}} \log(4+r^2) d\theta$$

$$= \frac{1}{2} \int_0^{\frac{\pi}{3}} (\log 8 - \log 4) d\theta = \frac{1}{2} (\log 8 - \log 4) \int_0^{\frac{\pi}{3}} d\theta$$

$$= \frac{1}{2} (\log 2) \int_0^{\frac{\pi}{3}} d\theta$$

$$\frac{\log 2}{2} \frac{\pi}{3} = \log 2 \left(\frac{\pi}{6} \right)$$

Answer: $\frac{\pi}{6} \log(2)$.