


Answer on Question #81220 Physics / Optics

Question. Newtons ring are formed with reflected light of wavelength 5895 \AA with a liquid between the plane and curved surface .The diameter of 5 *th* dark ring is 0.3 cm and radius of curvature is 1 m . Determine the refractive index of liquid?

Given. $\lambda = 5895 \text{ \AA} = 5895 \cdot 10^{-10} \text{ m}$; $m = 5$; $d_m = 0.3 \text{ cm} = 0.3 \cdot 10^{-2} \text{ m}$; $R = 1 \text{ m}$; $\beta = 0$.

Find. n —?

Solution.



For the dark ring system:

$$2dn \cos \beta + \frac{\lambda}{2} = (2m + 1)\lambda \rightarrow 2dn \cos \beta = m\lambda \rightarrow 2dn = m\lambda \rightarrow n = \frac{m\lambda}{2d}$$

From the figure

$$R^2 = (R - d)^2 + r^2$$
$$R^2 = R^2 - 2Rd + d^2 + r^2$$

Because d is very small, we have that

$$d = \frac{r^2}{2R}$$

Finally

$$n = \frac{4Rm\lambda}{d_m^2} = \frac{4 \cdot 1 \cdot 5 \cdot 5895 \cdot 10^{-10}}{(0.3 \cdot 10^{-2})^2} = 1.31$$

Answer. $n = 1.31$.

Answer provided by <https://www.AssignmentExpert.com>